

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

- 1. (Currently Amended) A method of prolonging a local effect of a pharmaceutically or cosmetically active substance when such substance is used in a topical cream or lotion prepared from an oil-in-water emulsion carrier, the method comprising:
- a. providing an oil-in-water emulsion carrier including an oily non-polar lipid material, an aqueous phase and a galactolipid material as an emulsifier;
- b. adding a pharmaceutically or cosmetically active substance to the carrier to form a formulation for a topical cream or lotion formulation;
- c. topically applying the topical cream or lotion formulation; and
- d. wherein the galactolipid material is present in the formulation in an amount to prolong the local effect of the pharmaceutically or cosmetically active substance when the





formulation is applied.

- 2. (Previously Amended) The method according to claim 1, wherein 0.1-50% by weight of oily material and 0.5-20% by weight of emulsifier are provided in the formulation.
- 3. (Previously Amended) The method according to claim 1, wherein 1-40% by weight of oily material and 0.5-10% by weight of emulsifier are provided in the formulation.
- 4. (Previously Amended) The method according to claim 1, including providing the galactolipid material so as to consist of at least 50% by weight of digalactosyl-diacylglycerols and a remainder of non-digalactosyldiacylglycerol polar lipids, and adding the galactolipid material so as to constitute an amount of 1.0-5.0% by weight of the formulation.
- 5. (Previously Amended) The method according to claim 1, including providing the galactolipid material so as to consist of 50-70% by weight of digalactosyldiacylglycerols and 30-50% by





weight of non-digalactosyldiacylglycerol polar lipids.

- 6. (Previously Amended) The method according to claim 1, including providing the galactolipid material as a fractionated oat oil which consists of at least 15% by weight of digalactosyldiacylglycerols and a remainder of non-digalactosyldiacylglycerol polar and non-polar lipids, and adding the galactolipid material so as to constitute an amount of 2.0-10% by weight of the formulation.
- 7. (Previously Amended) The method according to claim 6, including providing the galactolipid material as a fractionated oat oil which contains 40-60% by weight polar lipids and a remainder of non-polar lipids.
- 8. (Previously Amended) The method according to claim 1 wherein the oil-in-water emulsion is formed as a cream base by providing in % by weight;

Oily non-polar lipid

10.0-30.0%

. /t





Galactolipid emulsifier	0.5-5%
Thickner	2.0-10.0%
Preservative	0.1-1.0%
Water	ad 100%

- 9. (Previously Amended) The method according to claim 1 including incorporating a moisturizer as the active substance.
- 10. (Previously Amended) The method according to claim 1 including adding an active substance for use in treatment of atopic dermatitis.
- 11. (Previously Amended) The method according to claim 1 including incorporating a corticosteroid as the active substance, for treatment of skin inflammation.
- 12. (Previously Amended) The method according to claim 1 including incorporating 13-hydroxy-linoleic acid as the active substance.



13. (Previously Added) The method of claim 9 including adding glycerol as the moisturizer.